Quality of Life Improvement by Early Detection of Hearing Loss Using High-frequency Audiometry: Extended high-frequency audiometry as a screening method (Poster resume)

Michaela Skerkova¹, Martina Kovalova², Tomas Rychly³,Eva Mrazkova⁴, Hana Tomaskova⁵, Hana Slachtova⁶, Rastislav Madar⁷

¹MA, student of doctoral degree, Dept of Epidemiology and Public Health, Faculty of Medicine;
²MA, PhD, postdoctoral student, Dept of Epidemiology and Public Health;
³ Medical student;
^{1,2,3} Faculty of Medicine, University of Ostrava, Czech Republic
⁴ MD, PhD, researcher,
⁵ Ass. Prof. PhD; researcher,
⁶ MA, PhD, researcher;
⁷Ass. Prof, MD, PhD, FRCPS, Dean;
^{4,5,6,7} Dept. of Epidemiology and Public Health, Faculty of Medicine, University of Ostrava, Czech Republic;
^{5,6} Centre for Epidemiological Research, Faculty of Medicine, University of Ostrava, Czech Republic

First vision:

The aim of the project is to compare hearing loss using high-frequency audiometry, to find a fast, cheap and simple screening method for hearing testing in adults. The project will contribute to more effective prevention, will develop recommendations and guidelines for a hearing loss screening. In the first year of the project, 253 persons aged 18-65 years were examined, out of the sample 69% were women. The age groups were divided into 5 categories from 18-24 and further by 10 years of age. The preliminary results (the data collection has not been finished yet) shows higher hearing loss in men comparing with women and significantly worse results for right ear (p < 0.05) in both sexes. Statistically significant difference in high-frequency hearing loss between men and women was confirmed especially in the oldest age group (p < 0.05). When comparing differences of hearing loss between the frequency 9 kHz and 14 kHz significantly higher loss was confirmed for the frequency 14 kHz for all age groups and also right and left ear (p < 0.01). Based on the preliminary results, increased hearing threshold values were found in relationship with frequency, sex and age in unexposed population.

Recomended: Normative data for high-frequency audiometry have not yet been established. Therefore, further research will be necessary to establish a uniform evaluation protocol accounting for age, sex, comorbidities and exposures.

Keywords: High frequency audiometry; Hearing loss; General unexposed population.

Quality of Life Improvement by Early Detection of Hearing Loss Using High-frequency Audiometry: Extended high-frequency audiometry as a screening method (Poster resume)



Michaela Skerkova¹, Martina Kovalova², Tomas Rychly³,Eva Mrazkova⁴, Hana Tomaskova⁵, Hana Slachtova⁶, Rastislav Madar⁷

¹ MA, student of doctoral degree, Dept of Epidemiology and Public Health, Faculty of Medicine;
 ² MA, PhD, postdoctoral student, Dept of Epidemiology and Public Health; ³ Medical student;
 ^{1,2,3} Faculty of Medicine, University of Ostrava, Czech Republic; ⁴ MD, PhD, researcher;
 ⁵ Ass. Prof. PhD; researcher; ⁶ MA, PhD, researcher; ⁷Ass. Prof, MD, PhD, FRCPS, Dean;
 ^{45,6,7} Dept. of Epidemiology and Public Health, Faculty of Medicine, University of Ostrava, Czech Republic;
 ^{5,6} Centre for Epidemiological Research, Faculty of Medicine, University of Ostrava, Czech Republic

BACKGROUND AND AIM OF THE STUDY

The early detection and compensation of hearing deficiency enable full involvement of people into economic and social life.

The aim of the project is to compare hearing loss using high-frequency audiometry, to find a fast, cheap and simple screening method for hearing testing in adults. The project will contribute to more effective prevention, will develop recommendations and guidelines for a hearing loss screening.

RESULTS

In the first year of the project, 253 persons aged 18-65 years were examined, out of the sample 69% were women.

The age groups were divided into 5 categories from 18-24 and further by 10 years of age.

Figures 1-2 Hearing loss in men for right and left ear




The results shows higher hearing loss in men comparing with women and significantly worse results for right ear (p < 0.05) in both sexes.

Statistically significant difference in high-frequency hearing loss between men and women was confirmed especially in the oldest age group (p < 0.05). When comparing differences of hearing loss between the frequency 9 kHz and 14 kHz significantly higher loss was confirmed for the frequency 14 kHz for all age groups and also right and left ear (p < 0.01).

METHODS

The study of health status and hearing threshold has been provided using the cross-sectional design. The inclusion criteria: the population of both sexes in productive age, individual collection of the sample, the Czech nationality, absence of occupational exposure. Hearing threshold examinations consist of tympanometry, conventional pure tone audiometry of frequency 0.125-8 kHz, extended high-frequency audiometry of frequency 9-16 kHz, speech audiometry and transient evoked otoacoustic emissions. Hearing loss was evaluated based on medians. Comparison by sex age and frequency was analysed by Wilcoxon signed-rank test and Man-Whitney test on the significance level of 5% using the SW Stata v.14.





CONSLUSION

Based on the preliminary results, increased hearing threshold values were found in relationship with frequency, sex and age in unexposed population.

The results of this study will be used for determination of normative values for hearing loss in high frequencies as a guideline for assessment of hearing loss in workers exposed to noise.

Funding: Technology Agency of the Czech Rep. (TJ04000059) Contact details: michaela.skerkova@osu.cz 94

Caucasus Journal of Health Sciences and Public Health, Volume 5, Supplement 9